

## Treatment of Reperfusion Injury With Intracoronary Calcium Channel Antagonists and Reduced Coronary Free Calcium Concentration in Regionally Ischemic, Reperfused Porcine Hearts

HERMANN H. KLEIN, MD, SIBYLLE PICH, PhD, STEFANIE LINDERT,  
KLAUS NEBENDAHL, MD VET, GERHARD WARNEKE, MD,  
HEINRICH KREUZER, MD

Göttingen, West Germany

The effect of intracoronary diltiazem, EGTA (ethylene-bis-( $\beta$ -aminomethylether)-N,N'-tetraacetic acid), nifedipine, verapamil and isotonic saline solution as placebo on reperfusion injury was investigated in regionally ischemic, reperfused porcine hearts. The left anterior descending coronary artery was distally occluded for 45 min and was reperfused for 3 days. Intracoronary infusion was started immediately before reperfusion and continued during 45 min of reperfusion. Infarct size was determined as the ratio of infarcted (tetrazolium stain) to ischemic myocardium (dye technique). Regional systolic shortening was assessed by sonomicrometry.

Apart from left ventricular end-diastolic pressure before ischemia and during 45 min of reperfusion, global hemodynamic values in the five treatment groups did not differ; in particular, calculated left ventricular oxygen consumption before and during ischemia was equally low. Intracoronary EGTA decreased coronary venous free calcium concentration to about 70% of baseline value. Infarct size was reduced from  $76 \pm 10\%$  (control group,  $n = 8$ ) to  $60 \pm 10\%$  ( $p < 0.01$ ) by intracoronary diltiazem ( $n = 8$ ) and to  $55 \pm$

$15\%$  ( $p < 0.01$ ) by intracoronary EGTA ( $n = 8$ ). Insignificant reductions in infarct size were found after treatment with intracoronary verapamil ( $63 \pm 18\%$ ,  $n = 8$ ) and intracoronary nifedipine ( $68 \pm 9\%$ ,  $n = 7$ ). Regional systolic shortening of the risk region, which did not differ among the groups before occlusion and during ischemia, recovered to the greatest extent in the EGTA-treated pigs ( $p < 0.01$  compared with values in the control group). Treatment with intracoronary calcium antagonists resulted in only marginal improvement of systolic shortening. Reperfusion-induced rhythm disturbances were not beneficially affected by either treatment.

It is concluded that reperfusion injury exists in regionally ischemic, reperfused porcine hearts, which can be attenuated by intracoronary EGTA and intracoronary diltiazem administered during the first 45 min of reperfusion. The favorable action of EGTA is ascribed to a reduced ionized extracellular calcium concentration, whereas the beneficial mechanism of diltiazem is unknown.

(*J Am Coll Cardiol* 1989;13:1395-401)

Early restoration of blood flow represents a potent means to reduce infarct size and mortality in patients with acute myocardial infarction (1,2). Whether reperfusion of ischemic myocardium may cause cell death in myocytes that have survived the phase of ischemia is still subject to controversy. Treatments aimed at suppressing biochemical reactions that

are considered to be involved in the genesis of reperfusion injury in intact animals, e.g., scavenging of oxygen free radicals (3-6) or anti-inflammatory agents (7-9), have exhibited conflicting results. Sudden intracellular uptake of calcium during reperfusion (10), which can result in contraction band necrosis and mitochondrial accumulation of calcium (11), appears to be another mechanism for the evolution of reperfusion injury. In this study we examined whether intracoronary treatment with three calcium channel antagonists administered only during reperfusion, and lowered intracoronary free calcium concentration during reperfusion, would reduce infarct size and improve regional systolic shortening in porcine hearts subjected to 45 min of ischemia and 3 days of reperfusion.

From the Departments of Cardiology, Nephrology and Experimental Animal Research, University of Göttingen, Göttingen, West Germany. This study was supported by a grant from the Deutsche Forschungsgemeinschaft, SFB 330-Organprotektion, Göttingen.

Manuscript received August 20, 1988; revised manuscript received November 3, 1988, accepted December 7, 1988.

Address for reprints: Priv. Doz. Dr. H. H. Klein, Med. Klinik der Universität, Robert-Koch-Str. 40, 3400 Göttingen, West Germany.

## Methods

Medication, anesthesia, general surgical procedures and measurement of infarct size have been described in detail in previous studies (6,9,12). Fifty male and female farm pigs weighing 35 to 48 kg were used in this study.

**General experimental design.** After median thoracotomy, the left anterior descending coronary artery was dissected free at the beginning of its distal third. Left ventricular pressure and its first derivative ( $dp/dt$  max) were measured with a 5F Millar catheter-tipped manometer. Blood pressure was assessed in the ascending aorta with a fluid-filled catheter connected to a Statham transducer. A 5F multipurpose catheter was advanced from a jugular vein through the coronary sinus into the ostium of the great cardiac vein. A 4F Rentrop catheter with a 3F tip (USCI) was introduced under fluoroscopic control through the right carotid artery into the proximal part of the left anterior descending coronary artery. One pair of ultrasonic crystals (13) was implanted in the subendocardial layer of the planned ischemic region oriented parallel to the short axis (14).

**Regional systolic shortening** was determined by the ultrasonic transit time method (15,16). Analogue tracings of the signals were obtained from direct writing recordings. The end-diastolic distance of the crystals was determined at the onset of ventricular systole. Relative systolic shortening was assessed as the difference of distances between the beginning of ventricular systole and aortic valve closure over end-diastolic distance  $\times 100$ . Regional systolic shortening and global hemodynamic variables, including left ventricular peak pressure, left ventricular end-diastolic pressure, diastolic blood pressure,  $dp/dt$  max and heart rate, were recorded before occlusion, at 5 min intervals during ischemia and during 45 min of reperfusion and after 3 days of reperfusion. Left ventricular oxygen consumption ( $M\dot{V}O_2$ ) was calculated from the hemodynamic data before occlusion and during ischemia with use of Bretschneider's equation (17).

*The coronary artery was occluded at the prepared site for 45 min and was reperfused for 3 days.* In this preparation, occlusion of the coronary artery at the specified site renders about 15 g of the left ventricular muscle (average weight 120 g) ischemic. Forty-five minutes after the onset of reperfusion, the chest was closed in layers and the animal was allowed to recover. On the third postoperative day, the pig was reanesthetized and thoracotomy was repeated. After measurement of global hemodynamics and regional systolic shortening, the coronary artery was occluded at exactly the same site, a central venous injection of 20 ml 10% fluorescein sodium solution was administered to label the well perfused myocardium with a green fluorescent color and the heart was excised. The area at risk of necrosis was determined by planimetry of the nonfluorescent myocardium of four heart slices with use of enlarged photographs (13  $\times$  18 cm).

Infarcted myocardium was assessed by incubating the heart slices in nitroblue tetrazolium solution (18,19). Infarct size was calculated as a ratio of infarcted myocardium over risk region  $\times 100$ .

**Experimental protocol.** Fifty pigs were randomly assigned to one of the four treatment groups or to the control group. The intracoronary infusion of the active compounds or of isotonic sodium chloride solution as placebo was started 60 s before release of the coronary occlusion and was continued for 45 min during reperfusion. The four treatment groups, each consisting of 10 pigs, received either intracoronary diltiazem (Gödecke) (0.25 mg/min for 16 min followed by 0.125 mg/min for 30 min), intracoronary nifedipine (Bayer) (0.05 mg/min for 16 min followed by 0.025 mg/min for 30 min), intracoronary verapamil (Knoll) (0.125 mg/min for 16 min followed by 0.062 mg/min for 30 min) or intracoronary EGTA (ethylene-bis-( $\beta$ -aminomethylether)-N,N'-tetraacetic acid; analytical grade; Serva) neutralized to a pH value of 7.4 with sodium hydroxide. The dosage of EGTA was adjusted according to systolic blood pressure and ranged between 20 and 32  $\mu$ mol/min. In general, the initial dose was 28  $\mu$ mol EGTA/min, and the concentration was reduced when systolic blood pressure decreased to  $<70$  mm Hg.

*Coronary venous blood concentrations of total and free calcium* were determined with standard laboratory techniques before occlusion and after 2, 5, 10, 20 and 40 min of reperfusion. Arterial calcium concentrations were measured before and after 40 min of reperfusion.

**Statistics.** All data are presented as mean values  $\pm$  SD. Overall statistical comparisons among the groups were performed with the Kruskal-Wallis H test. When this test indicated statistical significance, the results of the different groups were analyzed with the Wilcoxon-Mann-Whitney U test (20). Results within a group were compared with the Wilcoxon test for paired observations. Probability values  $<5\%$  ( $p < 0.05$ ) were considered statistically significant.

## Results

During the first postoperative night, premature deaths occurred in 10 pigs—2 each in the diltiazem, verapamil and control groups, respectively, 3 in the nifedipine group and 1 in the EGTA group. In the latter group, one pig that developed ventricular fibrillation on reperfusion could not be resuscitated. Therefore, the results are based on 7 experiments in the nifedipine group and on 8 in each of the other four groups (total 39 pigs).

**Rhythm disturbances.** Ventricular fibrillation during ischemia occurred in 15 pigs: 2 in the diltiazem group, 3 each in the nifedipine, verapamil and control groups and 4 in the EGTA group. In most experiments, reperfusion induced various types of tachyarrhythmias, including accelerated idioventricular rhythms, premature ventricular complexes,

**Table 1.** Global Hemodynamic Variables In the Five Treatment Groups

Group	Before Occlusion	During Ischemia	Reperfusion	
			45 min	3 days
LVPP (mm Hg)				
D	99 ± 14	98 ± 13	91 ± 6	91 ± 17
E	93 ± 11	90 ± 12	79 ± 11*	82 ± 17
N	105 ± 18	100 ± 12	92 ± 9	101 ± 17
V	104 ± 15	100 ± 7	81 ± 9*	105 ± 18
C	107 ± 14	94 ± 15*	86 ± 13*	104 ± 6
LVEDP (mm Hg)				
D	13 ± 6	13 ± 5	12 ± 7	11 ± 4
E	6 ± 2	9 ± 2*	8 ± 3	10 ± 3
N	12 ± 6	15 ± 5	16 ± 5	13 ± 6
V	9 ± 4	9 ± 3	8 ± 4	10 ± 4
C	11 ± 5	10 ± 6	8 ± 3	12 ± 4
DBP (mm Hg)				
D	61 ± 15	64 ± 13	61 ± 9	61 ± 18
E	62 ± 10	60 ± 11	58 ± 9	51 ± 19
N	72 ± 20	71 ± 13	66 ± 9	70 ± 12
V	66 ± 9	66 ± 10	56 ± 11*	71 ± 18
C	74 ± 18	68 ± 14	64 ± 11	70 ± 10
Heart rate (beats × min <sup>-1</sup> )				
D	77 ± 25	86 ± 20*	106 ± 10*	91 ± 8
E	72 ± 9	74 ± 8	108 ± 14*	82 ± 17
N	72 ± 20	76 ± 14	106 ± 16*	92 ± 8*
V	61 ± 7	71 ± 10	102 ± 19*	90 ± 9*
C	70 ± 17	88 ± 14*	118 ± 14*	105 ± 14*
dP/dt max (mm Hg × sec <sup>-1</sup> )				
D	1,738 ± 534	1,720 ± 478	1,686 ± 450	1,162 ± 177*
E	1,775 ± 349	1,567 ± 218	1,283 ± 306*	1,088 ± 295*
N	2,028 ± 756	1,759 ± 510	1,745 ± 682	1,429 ± 330
V	1,650 ± 421	1,689 ± 468	1,377 ± 376*	1,200 ± 177*
C	1,662 ± 496	1,638 ± 583	1,639 ± 361	1,488 ± 461
MVO <sub>2</sub> (ml × min <sup>-1</sup> × 100 g <sup>-1</sup> )				
D	5.9 ± 1	6.3 ± 0.8*		
E	5.7 ± 0.7	5.6 ± 0.7		
N	5.9 ± 1.8	5.8 ± 0.8		
V	5.3 ± 0.5	5.7 ± 0.8		
C	5.6 ± 1.2	5.9 ± 0.8		


\*Significantly different compared with preischemic value. Values are mean ± SD. C = control; D = diltiazem; DBP = diastolic blood pressure; E = EGTA (ethylene-bis-(β-aminoethylether)-N,N'-tetraacetic acid); EDP = end-diastolic pressure; LV = left ventricular; MVO<sub>2</sub> = myocardial oxygen consumption; N = nifedipine; pp = peak systolic pressure; V = verapamil.

short runs of ventricular tachycardia and ventricular fibrillation. Ventricular fibrillation on reperfusion was observed in three pigs in the EGTA group, two in the verapamil group and one in the diltiazem group. It did not occur in the nifedipine-treated or control groups. Except for one pig in the EGTA group, ventricular fibrillation was always immediately reversed by direct electrical counter-shocks.

*Reperfusion-induced rhythm disturbances* were characterized by a sudden increase in heart rate. Comparison of the heart rates measured immediately before and after 1, 2, 3, 4, 5 and 10 min of reperfusion did not indicate any beneficial effect of the four treatments on reperfusion-induced tachy-

arrhythmias. The mean increases in heart rate were 24% (diltiazem group), 73% (EGTA group), 38% (nifedipine group), 37% (verapamil group) and 21% (control group).

**Global hemodynamics (Table 1).** Significant differences among the groups (Kruskal-Wallis test) were observed only for left ventricular end-diastolic pressure before occlusion and during 45 min of reperfusion. Before occlusion, left ventricular end-diastolic pressure was significantly lower in the EGTA group than in the nifedipine ( $p < 0.01$ ) or the diltiazem ( $p < 0.01$ ) group. During 45 min of reperfusion, it was significantly higher in the nifedipine-treated pigs than in the pigs in the control ( $p < 0.01$ ), EGTA ( $p < 0.01$ ) and verapamil-treated ( $p < 0.02$ ) groups. Calculated left ventric-




**Figure 1.** Free coronary venous calcium concentration in the EGTA-treated animals before treatment and during 40 min of reperfusion. The dosage of intracoronary (i.c.) EGTA (ethylene-bis-(β-aminoethylether)-N,N'-tetraacetic acid) is given at the bottom of the figure. Ca<sup>++</sup> = calcium ion; mval = mmol/valence.

ular oxygen consumption before and during ischemia was equally low in all groups.

**Effect of EGTA treatment on free coronary vein calcium concentration.** Coronary vein calcium concentrations did not change significantly in calcium antagonist and control experiments during reperfusion. Total arterial and coronary vein calcium concentrations that amounted to  $4.9 \pm 0.3$  mval/liter were not significantly affected by intracoronary EGTA treatment. Before occlusion, free coronary vein calcium concentrations were slightly ( $0.06 \pm 0.03$  mval/liter) but significantly higher than arterial concentrations. Intracoronary EGTA therapy reduced mean free coronary vein calcium concentrations to about 70% of baseline values (Fig. 1).

**Infarct size.** Risk regions of the five groups did not differ. The areas at risk of necrosis that were determined on photographs measured  $980 \pm 354$  mm<sup>2</sup> (diltiazem group),  $1,278 \pm 442$  mm<sup>2</sup> (EGTA group),  $1,218 \pm 336$  mm<sup>2</sup> (nifedipine group),  $1,043 \pm 170$  mm<sup>2</sup> (verapamil group) and  $1,184 \pm 246$  mm<sup>2</sup> (control group). Infarct size in the five groups is depicted in Figure 2. Treatment with diltiazem and with EGTA significantly reduced mean necrosis formation by 21% ( $p < 0.01$ ) and 28% ( $p < 0.01$ ), respectively. Infarct size in these two groups was also significantly smaller compared with that in the nifedipine treatment group ( $p < 0.05$ ). Reduction in infarct size by verapamil did not reach statistical significance, although mean infarct size in this group was only 5% larger than that in the group treated with diltiazem.


**Recovery of regional systolic shortening.** Baseline values of the end-diastolic distances of the ultrasonic crystals did not differ. These were  $10.6 \pm 2.8$  mm (diltiazem group),  $10.7 \pm 1.8$  mm (EGTA group),  $9.8 \pm 1.7$  mm (nifedipine



**Figure 2.** Infarct size of the five treatment groups after 45 min of ischemia and 3 days of reperfusion; p values are related to the infarct size of the control group. Other abbreviations as in Figure 1.

group),  $9.5 \pm 0.8$  mm (verapamil group) and  $10.4 \pm 2.4$  mm (control group). In all experiments, reperfusion induced a pronounced decrease in end-diastolic distance. After 10 min of reperfusion, mean end-diastolic distance was reduced to 75% (diltiazem group), 72% (EGTA group), 79% (nifedipine group), 74% (verapamil group) and 69% (control group) compared with the values immediately before reperfusion. Regional systolic shortening of the risk region in the five groups is shown in Figure 3. No difference among the groups was observed before ischemia and before reperfusion. After 3 days of reperfusion, regional systolic shortening had improved to the greatest extent in the EGTA-treated pigs ( $p <$

**Figure 3.** Regional systolic shortening of the risk region of the five groups; p values are related to regional systolic shortening of the control group after 3 days of reperfusion (reperf.). Other abbreviations as in Figure 1.



0.01 compared with control). Recovery of regional myocardial function of the diltiazem-, nifedipine- and verapamil-treated animals reached borderline significance.

## Discussion

**Calcium in regionally ischemic, reperfused myocardium.** Myocardium reversibly injured by 10 min of transient ischemia does not accumulate calcium. However, 40 min of regional ischemia followed by reperfusion results in an 18-fold increase of calcium in the damaged tissue (21), which occurs during the early phase of reperfusion (22). Much of the intracellular calcium is found in dense bodies within mitochondria. The process of mitochondrial uptake of calcium does not indicate that these organelles have died already. On the contrary, intact mitochondria prefer to accumulate calcium in preference to phosphorylation of adenosine triphosphate (ATP) (23). Mitochondria overloaded with calcium exhibit profound functional deterioration. Oxidative phosphorylating activity and ATP-generating capacity become considerably impaired (23,24). Whether the intracellular uptake of calcium during reperfusion determines the transition from reversible to irreversible injury in some jeopardized myocytes is still unresolved. Various studies (25-27) in globally ischemic heart preparations demonstrated that functional recovery after transient ischemia can be improved by a reduced extracellular calcium concentration. On the other hand, elevated extracellular calcium concentrations abolished functional impairment observed after short-term ischemia in regionally ischemic, reperfused canine hearts (28). Reperfusion of regionally ischemic canine myocardium with calcium-free blood revealed contradictory results in regard to ultrastructural cell damage. It was reported in one study (29) that this treatment preserved morphology after 45 min of transient ischemia, whereas another study (30) that used a similar protocol failed to demonstrate this beneficial effect.

*The results of the present study support the concept that reperfusion injury exists in regionally ischemic hearts.* Reduced free coronary calcium concentrations during 45 min of reperfusion significantly decreased infarct size and improved recovery of regional systolic shortening. Although reduced afterload has been found to favorably affect systolic bulging during ischemia (31), it is unlikely that the lower left ventricular peak pressure in the EGTA group accounts for the better recovery of regional function after 3 days of reperfusion. Whether EGTA treatment only accelerated functional recovery or whether this therapy induces ultimate improvement of systolic shortening by reducing infarct size cannot be answered with certainty. Because we were unable to measure free calcium concentrations in the arterial blood that perfused the risk region, we had to rely on blood concentrations determined in the corresponding vein. Extrapolation of coronary vein to coronary artery free calcium

concentration may result in a small error. Because free coronary vein calcium concentrations could not be measured on-line, it was impossible to achieve equal reductions of ionized calcium by the infusion of EGTA.

*Administration of intracoronary EGTA is not without risks.* Preliminary experiments had demonstrated that the therapeutic range of intracoronary EGTA is rather small. The intracoronary infusion of 40  $\mu$ mol EGTA/min, which decreased free coronary vein calcium concentration to about 0.6 mval/liter, induced ventricular tachycardia, and higher intracoronary EGTA doses always resulted in ventricular fibrillation. The optimal treatment durations and dosages of intracoronary EGTA and of intracoronary calcium antagonists during reperfusion are not known. We treated the pigs during 45 min of reperfusion because it was demonstrated that 20 min of reperfusion was sufficient to reverse ultrastructural changes of reversibly injured myocytes (32).

**Calcium antagonists for the treatment of reperfusion injury.** The mechanisms involved in the cellular uptake of calcium during reperfusion are not clearly defined. It is suggested that calcium enters the myocyte primarily by an increased permeability of the sarcolemma to calcium, which does not require membrane disruption (33,34). Whether calcium may also enter the cells by a sodium-calcium exchange (35,36) or by the voltage-dependent slow calcium channel is subject to controversy. Furthermore, the role of a fast calcium channel (37,38) for the cellular uptake of calcium on reperfusion has not yet been studied. In an isolated septum preparation, neither verapamil (39) nor nifedipine (40) was able to prevent intracellular calcium accumulation when the drug was administered during reperfusion. However, diltiazem was partially protective in a papillary muscle preparation (41) and nifedipine delayed the intracellular uptake of calcium in isolated rat hearts (34).

*At least 37 studies have been performed to test the effect of calcium antagonists on infarct size in different animal models (reviewed in Ref. 42).* In three of them (43-45), a calcium antagonist was administered only during ischemia and during reperfusion, which lasted for 3 to 4 h (43,44) or 1 week (45). Infarcted myocardium was delineated by tetrazolium staining. In two of the studies (43,44), infarct size was insignificantly reduced by either intracoronary verapamil (43) or intravenous diltiazem (44). However, a reperfusion period of 3 to 4 h, which was used in these studies, may have been too short to identify accurately necrotic myocardium by a tetrazolium stain when areas of no reflow were present (19). In the third study (45), nisoldipine, a derivative of nifedipine, was ineffective in treating reperfusion injury in baboons. In that study and in the canine study (43) that tested verapamil, reperfusion was started only after 3 h of ischemia, a time when mitochondrial accumulation of calcium may no longer occur (11). In the present study, the dosages of the calcium antagonists were guided by therapeutic plasma levels of these compounds in humans (46). The

intracoronary infusion of the calcium antagonists warranted blood concentrations in the coronary artery that were  $\geq 2.5$ - to 5-fold higher than usual therapeutic plasma levels, even when maximal coronary vasodilation was taken into account.

**Comparison of the three calcium antagonists.** The reason for the different effects of the three calcium antagonists on infarct size remains to be elucidated. Although the vasodilator adenosine administered during reperfusion has reduced infarct size and improved functional recovery in canine hearts (47), it is unlikely that the results of our study can be ascribed to different degrees of vasodilation because nifedipine is a potent vasodilator (48). The three calcium antagonists produced marginal improvement in regional systolic shortening. Whether this effect can be extrapolated to a better ultimate improvement of regional function due to infarct size reduction is uncertain because it has been demonstrated (49) that verapamil attenuated postischemic dysfunction in canine hearts after short-term ischemia when no infarctions were present.

**Conclusions.** Our study suggests that reperfusion injury exists in regionally ischemic porcine hearts and that this injury can be attenuated by intracoronary EGTA and intracoronary diltiazem administered during 45 min of reperfusion. At the given drug dosages, diltiazem was superior to nifedipine but only insignificantly better than verapamil.

## References

1. The I.S.A.M. Study Group. A prospective trial of intravenous streptokinase in acute myocardial infarction (ISAM). *N Engl J Med* 1986;314:1465-71.
2. Gruppo Italiano per lo Studio della Streptochinasi nell'Infarto Miocardico (GISSI). Effectiveness of intravenous thrombolytic treatment in acute myocardial infarction. *Lancet* 1986;1:397-401.
3. Werns SW, Shea MJ, Driscoll EM, et al. The independent effects of oxygen radical scavengers on canine infarct size. Reduction by superoxide dismutase but not catalase. *Circ Res* 1985;56:895-8.
4. Ambrosio G, Becker LC, Hutchins GM, Weisman HF, Weisfeld ML. Reduction in experimental infarct size by recombinant human superoxide dismutase: insights into the pathophysiology of reperfusion injury. *Circulation* 1986;74:1424-33.
5. Uraizee A, Reimer KA, Murry C, Jennings RB. Failure of superoxide dismutase to limit size of myocardial infarction after 40 minutes of ischemia and 4 days of reperfusion in dogs. *Circulation* 1987;75:1237-48.
6. Klein HH, Pich S, Lindert S, Buchwald A, Nebendahl K, Kreuzer H. Intracoronary superoxide dismutase for the treatment of reperfusion injury: a blind randomized placebo-controlled trial in ischemic, reperfused porcine hearts. *Basic Res Cardiol* 1988;83:141-8.
7. Jolly SR, Lucchesi BR. Effect of BW755C in an occlusion-reperfusion model of ischemic myocardial injury. *Am Heart J* 1983;106:8-13.
8. Mullane KM, Read N, Salmon JA, Moncada S. Role of leukocytes in acute myocardial infarction in anesthetized dogs: relationship to myocardial salvage by anti-inflammatory drugs. *J Pharmacol Exp Ther* 1984;228:510-22.
9. Klein HH, Pich S, Bohle RM, et al. The anti-inflammatory agent BW755C in ischemic, reperfused porcine hearts. *J Cardiovasc Pharmacol* 1988;12:338-44.
10. Shen AC, Jennings RB. Myocardial calcium and magnesium in acute ischemic injury. *Am J Pathol* 1972;67:417-40.
11. Jennings RB. Calcium ions in ischemia. In: Opie LH, ed. *Calcium Antagonists and Cardiovascular Disease*. New York: Raven, 1984:85-95.
12. Pich S, Klein HH, Lindert S, Nebendahl K, Buchwald A, Kreuzer H. Cell death in ischemic, reperfused porcine hearts: a histochemical and functional study. *Basic Res Cardiol* 1988;83:550-9.
13. Heimisch W, Hagl S, Gebhardt K, Meisner H, Mandler N, Sebening F. Direct measurement of cyclic changes in regional wall geometry in the left ventricle of the dog. *Innov Tech Biol Med* 1981;2:487-501.
14. Freeman GL, Le Winter MM, Engler RL, Covell JW. Relationship between myocardial fiber direction and segment shortening in the midwall of the canine left ventricle. *Circ Res* 1985;56:31-9.
15. Bugge-Asperheim B, Leraand S, Kill F. Local dimensional changes of the myocardium measured by ultrasonic technique. *Scand J Clin Lab Invest* 1969;24:361-71.
16. Theroux P, Franklin D, Ross J Jr, Kemper WS. Regional myocardial function during acute coronary artery occlusion and its modification by pharmacologic agents in the dog. *Circ Res* 1974;35:896-908.
17. Bretschneider HJ, Hellige G. Pathophysiologie der Ventrikelkontraktion-Kontraktilität, Inotropie, Suffizienzgrad und Arbeitsökonomie des Herzens. *Verh Dtsch Ges Kreislaufforsch* 1976;41:14-30.
18. Nachlas MU, Shnitka TK. Macroscopic identification of early myocardial infarcts by alteration in dehydrogenase activity. *Am J Pathol* 1963;42:379-405.
19. Klein HH, Puschmann S, Schaper J, Schaper W. The mechanism of the tetrazolium reaction in identifying experimental myocardial infarction. *Virchows Arch [A]* 1981;393:287-97.
20. Clauss G, Ebner H. Statistik. Thun, Frankfurt: Harri Deutsch Verlag, 1982:345-9.
21. Shen AC, Jennings RB. Kinetics of calcium accumulation in acute myocardial injury. *Am J Pathol* 1972;67:441-52.
22. Whalen DA Jr, Hamilton DG, Ganote G, Jennings RB. Effect of a transient period of ischemia on myocardial cells. I. Effects on cell volume regulation. *Am J Pathol* 1974;74:381-98.
23. Lehninger AL. Mitochondria and calcium ion transport. *Biochem J* 1970;119:129-38.
24. Nayler WG. The role of calcium in the ischemic myocardium. *Am J Pathol* 1981;102:262-70.
25. Follette DM, Fey K, Buckberg GD, et al. Reducing postischemic damage by temporary modification of reperfusate calcium, potassium, pH, and osmolality. *J Thorac Cardiovasc Surg* 1981;82:221-38.
26. Shine KI, Douglas AM. Low calcium reperfusion of ischemic myocardium. *J Mol Cell Cardiol* 1983;15:251-60.
27. Kuroda H, Ishiguro S, Mori T. Optimal calcium concentration in the initial reperfusate for post-ischemic myocardial performance (calcium concentration during reperfusion). *J Mol Cell Cardiol* 1986;18:625-33.
28. Ito BR, Tate H, Kobayashi M, Schaper W. Reversibly injured, postischemic canine myocardium retains normal contractile reserve. *Circ Res* 1987;61:834-46.
29. Ashraf M, White F, Bloor CM. Ultrastructural influences of reperfusing dog myocardium with calcium-free blood after coronary artery occlusion. *Am J Pathol* 1978;90:423-34.
30. Rocco MB, Reimer KA, Jennings RB. Failure of calcium chelated blood to prevent reperfusion induced indices of irreversible ischemic injury (abstr). *Circulation* 1979;59(suppl II):II-216.
31. Noma S, Askenase AD, Agarwalla JB, Helfant RH. The effect of changes in afterload on systolic bulging. *Circulation* 1988;77:221-6.
32. Jennings RB, Schaper J, Hill ML, Steenbergen C Jr, Reimer KA. Effect of reperfusion late in the phase of reversible ischemic injury. *Circ Res* 1985;56:262-78.
33. Crake T, Poole-Wilson PA. Evidence that calcium influx on reoxygenation is not due to cell membrane disruption in the isolated rabbit heart. *J Mol Cell Cardiol* 1986;18(suppl 4):31-6.

34. Nayler WG, Panagiotopoulos S, Elz JS, Daly MJ. Calcium-mediated damage during post-ischaemic reperfusion. *J Mol Cell Cardiol* 1988; 20(suppl II):41-54.
35. Grinwald PM, Brosnahan C. Sodium imbalance as a cause of calcium overload in post-hypoxic reoxygenation injury. *J Mol Cell Cardiol* 1987; 19:487-95.
36. Poole-Wilson PA, Tones MA. Sodium exchange during hypoxia and on reoxygenation in the isolated rabbit heart. *J Mol Cell Cardiol* 1988; 20(suppl II):15-22.
37. Bean BP. Two kinds of calcium channels in canine atrial cells: differences in kinetics, selectivity, and pharmacology. *J Gen Physiol* 1985;86:1-30.
38. Nilius B, Hess P, Lansman JB, Tsien RW. A novel type of cardiac calcium channel in ventricular cells. *Nature* 1985;316:443-6.
39. Bourdillon PD, Poole-Wilson PA. The effects of verapamil, quiescence, and cardioplegia on calcium exchange and mechanical function in ischemic rabbit myocardium. *Circ Res* 1982;50:360-8.
40. Poole-Wilson PA, Harding DP, Bourdillon PDV, Tones MA. Calcium out of control. *J Mol Cell Cardiol* 1984;16:175-87.
41. Chapell SP, Lewis MJ, Henderson AH. Myocardial reoxygenation damage: can it be circumvented. *Cardiovasc Res* 1985;19:299-305.
42. Kloner RA, Braunwald E. Effects of calcium antagonists on infarcting myocardium. *Am J Cardiol* 1987;59:84B-94B.
43. Lo H-M, Kloner RA, Braunwald E. Effect of intracoronary verapamil on infarct size in the ischemic, reperfused canine heart: critical importance of the timing of treatment. *Am J Cardiol* 1985;56:672-7.
44. Klein HH, Schuboth M, Nebendahl K, Kreuzer H. The effect of two different diltiazem treatments on infarct size in ischemic, reperfused porcine hearts. *Circulation* 1984;69:1000-5.
45. Vatner SF, Patrick TA, Knight DR, Manders WT, Fallon JT. Effects of calcium channel blocker on responses of blood flow, function, arrhythmias, and the extent of infarction following reperfusion in conscious baboons. *Cir Res* 1988;62:105-15.
46. Hugenoltz PG. Calcium antagonists. In: Abshagen U, ed. *Clinical Pharmacology of Antianginal Drugs*. Berlin, New York, Tokyo: Springer, 1985:459-538.
47. Olafsson B, Forman MB, Puett DW, et al. Reduction of reperfusion injury in the canine preparation by intracoronary adenosine: importance of the endothelium and the no-reflow phenomenon. *Circulation* 1987;76:1135-45.
48. Henry PD. Comparative pharmacology of calcium antagonists: nifedipine, verapamil, and diltiazem. *Am J Cardiol* 1980;46:1047-58.
49. Przyklenk K, Kloner PA. Effect of verapamil on postischemic "stunned" myocardium: importance of the timing of treatment. *J Am Coll Cardiol* 1988;11:614-23.